

SUBGROUPS AND QUOTIENTS OF AUTOMORPHISM GROUPS OF RAAGS

RUTH CHARNEY AND KAREN VOGTMANN

ABSTRACT. We study subgroups and quotients of outer automorphism groups of right-angled Artin groups (RAAGs). We prove that for all RAAGs, the outer automorphism group is residually finite and, for a large class of RAAGs, it satisfies the Tits alternative. We also investigate which of these automorphism groups contain non-abelian solvable subgroups.

1. INTRODUCTION

A right-angled Artin group, or RAAG, is a finitely-generated group determined completely by the relations that some of the generators commute. A RAAG is often described by giving a simplicial graph Γ with one vertex for each generator and one edge for each pair of commuting generators. RAAGs include free groups (none of the generators commute) and free abelian groups (all of the generators commute). Subgroups of free groups and free abelian groups are easily classified and understood, but subgroups of right-angled Artin groups lying between these two extremes have proved to be a rich source of examples and counterexamples in geometric group theory. For details of this history, we refer to the article [Ch07].

Automorphism groups and outer automorphism groups of RAAGs have received less attention than the groups themselves, with the notable exception of the two extreme examples, i.e. the groups $Out(F_n)$ of outer automorphism groups of a free group and the general linear group $GL(n, \mathbb{Z})$. The group $Out(F_n)$ has been shown to share a large number of properties with $GL(n, \mathbb{Z})$, including several kinds of finiteness properties and the Tits alternative for subgroups. These groups have also been shown to differ in significant ways, including the classification of solvable subgroups. In a series of recent papers [CCV07, CV08, BCV09], we have begun to address the question of which properties shared by $Out(F_n)$ and $GL(n, \mathbb{Z})$ are in fact shared by the entire class of outer automorphism groups of right-angled Artin groups. We are also interested in the question of determining properties which depend on the shape of Γ and in determining exactly how they depend on it.

In our previous work, an important role was played by certain restriction and projection homomorphisms, which allow one to reduce questions about the full outer automorphism group of a RAAG to questions about the outer automorphism groups of smaller subgroups.

R. Charney was partially supported by NSF grant DMS 0705396. K. Vogtmann was partially supported by NSF grant DMS 0705960.

In the first section of this paper we recall these tools and develop them further. In the next section we apply them to prove

Theorem 4.2. *For any defining graph Γ , the group $Out(A_\Gamma)$ is residually finite.*

This result was obtained independently by A. Minasyan [Mi09], by different methods. We next prove the Tits' alternative for a certain class of *homogeneous* RAAGs (see section 5).

Theorem 5.5. *If Γ is homogeneous, then $Out(A_\Gamma)$ satisfies the Tits' alternative.*

In the last section, we investigate solvable subgroups of $Out(A_\Gamma)$. We provide examples of non-abelian solvable subgroups and we determine an upper bound on the virtual derived length of solvable subgroups when A_Γ is homogeneous. Finally, by studying translation lengths of infinite order elements, we find conditions under which all solvable subgroups of $Out(A_\Gamma)$ are abelian. We show that excluding "adjacent transvections" from the generating set of $Out(A_\Gamma)$ gives rise to a subgroup $\tilde{Out}(A_\Gamma)$ satisfying a strong version of the Tits alternative.

Corollary 6.12. *If Γ is homogeneous of dimension n , then every subgroup of $\tilde{Out}(A_\Gamma)$ is either virtually abelian or contains a non-abelian free group.*

Thus for graphs which do not admit adjacent transvections, the whole group $Out(A_\Gamma)$ satisfies this property. One case which is simple to state is the following.

Corollary 6.13. *If Γ is connected with no triangles and no leaves, then all solvable subgroups of $Out(A_\Gamma)$ are virtually abelian.*

The authors would like to thank Talia Fernós for helpful conversations.

2. SOME COMBINATORICS OF SIMPLICIAL GRAPHS

Certain combinatorial features of the defining graphs Γ for our right-angled Artin groups will be important for studying their automorphisms. In this section we establish notation and recall some basic properties of these features.

Definition 2.1. Let v be a vertex of Γ . The *link* of v , denoted $lk(v)$, is the full subgraph spanned by all vertices adjacent to v . The *star* of v , denoted $st(v)$, is the full subgraph spanned by v and $lk(v)$.

Definition 2.2. Let Θ be a subgraph of Γ . The *link* of Θ , denoted $lk(\Theta)$, is the intersection of the links of all vertices in Θ . The *star* of Θ , denoted $st(\Theta)$ is the full subgraph spanned by $lk(\Theta)$ and Θ . The *perp* of Θ , denoted Θ^\perp , is the intersection of the stars of all vertices in Θ .

These can be expressed in terms of distance in the graph as follows:

- $v \in lk(\Theta)$ iff $d(v, w) = 1$ for all $w \in \Theta$
- $v \in \Theta^\perp$ iff $d(v, w) \leq 1$ for all $w \in \Theta$
- $v \in st(\Theta)$ iff $v \in lk(\Theta) \cup \Theta$

Recall that a complete subgraph of Γ is called a *clique*. (In this paper, cliques need not be maximal.) If Δ is a clique, then $st(\Delta) = \Delta^\perp$; otherwise $st(\Delta)$ strictly contains Δ^\perp .

Lemma 2.3. *If Δ is a clique, then $st(\Delta)^\perp$ is also a clique and $st(\Delta) \supseteq st(\Delta)^\perp \supseteq \Delta$.*

Proof. Since Δ is a clique, $v \in st(\Delta)$ implies $st(v) \supseteq \Delta$. Therefore

$$st(\Delta)^\perp = \bigcap_{v \in st(\Delta)} st(v) \supseteq \Delta.$$

If $x \in st(\Delta)^\perp$, then $d(x, v) \leq 1$ for all vertices $v \in st(\Delta)$, including all $v \in \Delta$, i.e. $x \in st(\Delta)$. If y is another vertex in $st(\Delta)^\perp$, then similarly $d(y, v) \leq 1$ for all vertices $v \in st(\Delta)$, so in particular $d(y, x) = 1$. Since any two vertices of $st(\Delta)^\perp$ are adjacent, $st(\Delta)^\perp$ is a clique. □

We define $v \leq w$ to mean $lk(v) \subseteq st(w)$. This relation is transitive and induces a partial ordering on equivalence classes of vertices $[v]$, where $w \in [v]$ if and only if $v \leq w$ and $w \leq v$ ([CV08], Lemma 2.2). The links $lk[v]$ and stars $st[v]$ of equivalence classes of maximal vertices v will be of particular interest to us.

The equivalence classes of maximal vertices of Γ form the vertices of an associated graph Γ_0 . There is an edge from $[v]$ to $[w]$ in Γ_0 if and only if there is an edge from v to w in Γ . The graph Γ_0 can be realized as a subgraph of Γ and if Γ is connected, then so is Γ_0 .

For a full subgraph $\Theta \subset \Gamma$, the right-angled Artin group A_Θ embeds into A_Γ in the natural way. The image is called a *special subgroup* of A_Γ , and we use the same notation A_Θ for it. An important observation is that the centralizer of A_Θ is equal to A_{Θ^\perp} (see, e.g., [CCV07], Proposition 2.2).

We remark that if v is a vertex in $\Theta \subset \Gamma$, then it is possible for v to be maximal in Θ but not in Γ . Unless otherwise stated, the term "maximal vertex" will always mean maximal with respect to the original graph Γ .

The subgraph spanned by $[v]$ is either a clique, or it is disconnected and discrete ([CV08], Lemma 2.3). In the first case the subgroup $A_{[v]}$ is abelian and we call v an *abelian vertex*; in the second, $A_{[v]}$ is a non-abelian free group, and we call v a *non-abelian vertex*. Note that for any vertex v , $st[v]$ is the union of the stars of the vertices $w \in [v]$.

A *leaf* of Γ is a vertex which is an endpoint of only one edge. A *leaf-like* vertex is a vertex v whose link contains a unique maximal vertex w , and $[v] \leq [w]$. In particular, a leaf is leaf-like. If Γ has no triangles, then every leaf-like vertex is in fact a leaf.

3. KEY TOOLS

Generators for $Out(A_\Gamma)$ were determined by M. Laurence [Lau95], extending work of H. Servatius [Ser89]. They consist of

- graph automorphisms
- inversions of a single generator v
- transvections $v \mapsto vw$ for generators $v \leq w$
- partial conjugations by a generator v on one component of $\Gamma - st(v)$

As in [CV08], we consider the finite-index subgroup $Out^0(A_\Gamma)$ of $Out(A_\Gamma)$ generated by inversions, transvections and partial conjugations. This is a normal subgroup, called the *pure outer automorphism group*

If Γ is connected and v is a maximal vertex, then any pure outer automorphism ϕ of A_Γ has a representative f_v which preserves both $A_{[v]}$ and $A_{st[v]}$ ([CV08], Prop. 3.2). This allows us to define several maps from $Out^0(A_\Gamma)$ to the outer automorphism groups of various special subgroups, as follows.

- (1) Restricting f_v to $A_{st[v]}$ gives a *restriction map*

$$R_v: Out^0(A_\Gamma) \rightarrow Out^0(A_{st[v]}).$$

- (2) The map $A_\Gamma \rightarrow A_{\Gamma-[v]}$ which sends each generator in $[v]$ to the identity induces an *exclusion map*

$$E_v: Out^0(A_\Gamma) \rightarrow Out^0(A_{\Gamma-[v]}).$$

- (3) Since v is maximal with respect to the graph $st[v]$ and $lk[v] = st[v] - [v]$, we can compose the restriction map on A_Γ with the exclusion map on $A_{st[v]}$ to get a *projection map*

$$P_v: Out^0(A_\Gamma) \rightarrow Out^0(A_{lk[v]}).$$

If Γ is the star of a single vertex v , then $[v]$ is the unique maximal equivalence class, and R_v is the identity. If Γ is a complete graph, then $\Gamma = [v]$ and $lk[v]$ is empty, in which case we define $P_v = E_v$ to be the trivial map.

3.1. The amalgamated restriction homomorphism R . Let Γ be a connected graph. We can put all of the restriction maps R_v together to obtain an amalgamated restriction map

$$R = \prod R_v: Out^0(A_\Gamma) \rightarrow \prod Out^0(A_{st[v]}),$$

where the product is over all maximal equivalence classes $[v]$. It was proved in [CV08] that the kernel K_R of R is a finitely-generated free abelian group, generated by partial conjugations. If Γ has no triangles, we also found a set of generators for K_R [CCV07]. We will need this information for general Γ in what follows, so we will now present another (and simpler) proof that K_R is free abelian which also identifies a set of generators for K_R .

By definition, any automorphism representing an element of K_R acts on the star of each maximal equivalence class of vertices as conjugation by some element of A_Γ . We begin by showing that the same is true for every equivalence class:

Lemma 3.1. *Let f be an automorphism representing an element of K_R . Then for every vertex $v \in \Gamma$, f acts on $st[v]$ as conjugation by some $g \in A_\Gamma$.*

Proof. This is by definition of the kernel if v is maximal. Since every vertex of Γ is in the star of some maximal vertex, f sends every vertex to a conjugate of itself. By a result of Laurence ([Lau95], Theorem 2.2), this implies that f is a product of partial conjugations.

If v is not maximal, then choose a maximal vertex v_0 with $v < v_0$. After adjusting by an inner automorphism if necessary, we may assume f is the identity on $st[v_0]$. If v is adjacent to v_0 , then $st[v] \subset st[v_0]$ and we are done.

If v is not adjacent to v_0 , choose a maximal vertex $w_0 \in lk(v) \cap lk(v_0)$ (note that one always exists). Then f acts as conjugation by some g on $st[w_0]$. Let e_0 be the edge from v_0 to w_0 . Since $st(e_0) \subset st(w_0)$, f acts as conjugation by g on all of $st(e_0)$. Since

$st(e_0) \subset st(v_0)$, g centralizes $st(e_0)$, i.e. g is in the subgroup generated by $st(e_0)^\perp$. By Lemma 2.3, $st(e_0)^\perp = \Delta$ is a clique containing e_0 , so the subgroup A_Δ is abelian.

Since A_Δ is abelian, we can write $g = g_2 g_1$ where g_1 is a product of generators in $lk[v]$ and g_2 a product of generators not in $lk[v]$. We claim that f acts as conjugation by g_2 on all of $st[v]$. Since $[v] \subset st[w_0]$, f acts as conjugation by g on $[v]$, and since g_1 commutes with $[v]$, this is the same as conjugation by g_2 . The action of f on $lk[v]$ is trivial, since $lk[v] \subset st[v_0]$, so it suffices to show that g_2 commutes with $lk[v]$. For suppose $u \in \Delta$ does not lie in $lk[v]$, and $x \in lk[v]$. Then either x lies in $st(u)$, hence commutes with u , or x and v lie in the same component of $\Gamma - st(u)$. In the latter case, since f is a product of partial conjugations, the total exponent of u in the conjugating element must be the same at v and at x ; but $f(x) = x$, so this total exponent must be 0. That is, u can appear as a factor in g_2 only if it commutes with all of $lk[v]$. \square

Next, we describe some automorphisms contained in the kernel K_R . If Γ is a connected graph and v is a vertex of Γ , say vertices x and y are in the same \hat{v} -component of Γ if x and y can be connected by an edge-path which contains no edges of $st(v)$ (though it may contain vertices of $lk(v)$). A \hat{v} -component lying entirely inside $st(v)$ is called a *trivial* \hat{v} -component, and any other \hat{v} -component is *non-trivial*. If $st(v)$ has no triangles, a non-trivial \hat{v} -component is the same thing as a non-leaf component of $\Gamma - v$. In general, each component of $\Gamma - st(v)$ is contained in a single \hat{v} -component, but a single \hat{v} -component may contain several components of $\Gamma - st(v)$.

Definition 3.2. A \hat{v} -component conjugation is an automorphism of A_Γ which conjugates all vertices in a single nontrivial \hat{v} -component of Γ by v .

By the remarks above, a \hat{v} -component conjugation is in general a product of partial conjugations by v on components of $\Gamma - st(v)$. To see that such conjugations lie in K_R , note that for any w , all of the vertices of $st[w]$ which do not lie in $st(v)$ lie in the same \hat{v} -component as w . Hence any \hat{v} -component conjugation acts as an inner automorphism on $st[w]$.

Let $\hat{c}(v)$ be the number of non-trivial \hat{v} -components in Γ .

Theorem 3.3. *The kernel K_R of the restriction map is free abelian, generated by non-trivial \hat{v} -component conjugations for all $v \in \Gamma$. The rank of K_R is $\sum_{v \in \Gamma} (\hat{c}(v) - 1)$.*

Proof. Let $\widehat{\mathcal{PC}}$ denote the set of all non-trivial \hat{v} -component conjugations for all $v \in \Gamma$. We first prove that $\widehat{\mathcal{PC}}$ generates K_R .

Let $\phi \in K_R$. For each representative f of ϕ , let V_f be the set of vertices v such that $st[v]$ is pointwise fixed. Now fix a representative f such that V_f is of maximal size. We proceed by induction on the number of vertices in $\Gamma - V_f$.

If $V_f = \Gamma$, then f is the trivial automorphism and there is nothing to prove. If not, choose a vertex w at distance 1 from V_f , so w is connected by an edge e to some $v \in V_f$. Then f acts non-trivially on $st[w]$ as conjugation by some $g \in A_\Gamma$. Since f acts trivially on $st[v]$, g fixes $st(e)$. The centralizer of $A_{st(e)}$ is equal to $A_{st(e)^\perp}$. By Lemma 2.3, $st(e)^\perp = \Delta$

for some clique Δ containing e , so g is in the abelian subgroup A_Δ and we can write $g = u_1^{\epsilon_1} \dots u_k^{\epsilon_k}$ for distinct vertices $u_i \in \Delta$.

If $st(w) \subseteq st(u_i)$ (e.g. if $u_i = w$), conjugation by u_i is trivial on $st(w)$ and we may assume $\epsilon_i = 0$, i.e. u_i does not appear in the expression for g . If $V_f \subset st(u_i)$, replace f by f composed with the inner automorphism by $u_i^{-\epsilon_i}$; the new V_f contains (so is equal to) the old one. We may now assume neither $st(w)$ nor V_f are contained in the star of any u_i .

Fix u_i and $x \in st(w) - st(u_i)$ and $y \in V_f - st(u_i)$. We claim that x and y are in different components of $\Gamma - st(u_i)$. To see this, suppose x and y are in the same component of $\Gamma - st(u_i)$. Since f sends each vertex to a conjugate of itself, f is a product of partial conjugations and x and y must be conjugated by the same total power of u_i . For y this power is zero, since $y \in V_f$, and so ϵ_i must also be zero, i.e. u_i does not occur in the expression for g .

We claim further that x and y must be in different \hat{u}_i -components of Γ . Suppose they were in the same \hat{u}_i -component. Let γ be an edge-path joining y to x which avoids edges of $st(u_i)$, with vertices $y = x_0, x_1, x_2, \dots, x_k = x$. We know that y is fixed by f and x is conjugated by a non-trivial power of u_i . Therefore there is some x_j in $lk(u_i)$ with the property that x_{j-1} is not conjugated by u_i but x_{j+1} is conjugated by a non-trivial power of u_i . Since γ does not use edges of $st(u_i)$, neither x_{j-1} nor x_{j+1} is in $lk(u_i)$, i.e. neither commutes with u_i . Thus f does not act as conjugation by the same total power of u_i on all of $st[x_i]$, contradicting Lemma 3.1.

The vertices of $st(w) - st(u_i)$ lie in a single, non-trivial \hat{u}_i -component (the component containing w) and by the discussion above, this component contains no vertices of $V_f - st(u_i)$. Thus, there is a non-trivial \hat{u}_i -component conjugation f_i which affects vertices of $st(w)$ but not V_f . The automorphism $f' = f_k^{-\epsilon_k} \circ \dots \circ f_1^{-\epsilon_1} \circ f$ has a strictly larger $V_{f'}$, which includes w as well as V_f . By induction, f' is a product of elements of $\widehat{\mathcal{PC}}$, hence so is f .

It remains to check that any two elements of $\widehat{\mathcal{PC}}$ commute in $Out(A_\Gamma)$. Let f_v be \hat{v} -component conjugation on C_v , and f_w a \hat{w} -component conjugation on C_w . If v and w are adjacent, these commute. If $d(v, w) > 1$, then $st(w)$ is contained in a single \hat{v} -component D_v , and $st(v)$ is contained in a single \hat{w} -component D_w . It follows that D_v contains every \hat{w} -component except D_w , and D_w contains every \hat{v} -component except D_v . It is now easy to check that any two components C_w and C_v are either disjoint, $C_v \subset C_w$, $C_w \subset C_v$ or $\Gamma - C_v$ and $\Gamma - C_w$ are disjoint; in any of these cases, the outer automorphisms associated f_v and f_w commute.

The only other relation among the generators of $\widehat{\mathcal{PC}}$ is that for a fixed v , the product of all non-trivial \hat{v} -component conjugations is an inner automorphism. The last statement of the theorem follows. \square

3.2. The amalgamated projection homomorphism P . We can combine the projection homomorphisms P_v for maximal equivalence classes $[v]$ in the same way we combined the restriction homomorphisms, to obtain an amalgamated projection homomorphism

$$P = \prod P_v : Out^0(A_\Gamma) \rightarrow \prod Out^0(A_{lk[v]}).$$

Recall that a vertex v is called leaf-like if there is a unique maximal vertex w in $lk(v)$ and this vertex satisfies $[v] \leq [w]$. The transvection $v \mapsto vw$ is called a *leaf transvection*. It is proved in [CV08] that the kernel K_P of P is a free abelian group generated by K_R and the set of all leaf transvections.

4. RESIDUAL FINITENESS

It is easy to see using congruence subgroups that $GL(n, \mathbb{Z})$ is residually finite, and E. Grossman proved that $Out(F_n)$ is also residually finite ([Gr74]). In this section we use these facts together with our restriction and exclusion homomorphisms to show that in fact $Out(A_\Gamma)$ is residually finite for every defining graph Γ . The same result has been obtained by A. Minasyan [Mi09] by different methods. Both proofs use a fundamental result of Minasyan and Osin which takes care of the case when the defining graph is disconnected:

Theorem 4.1. [MiOs09] *If G is a finitely generated, residually finite group with infinitely many ends, then $Out(G)$ is residually finite.*

Theorem 4.2. *For any right-angled Artin group A_Γ , $Out(A_\Gamma)$ is residually finite.*

Proof. Every right angled Artin group A_Γ is finitely generated and residually finite (it's linear), and A_Γ has infinitely many ends if and only if Γ is disconnected. Therefore, by Theorem 4.1, we may assume that Γ is connected.

We proceed by induction on the number of vertices in Γ .

Consider first the case in which $\Gamma = st[v]$ for a single equivalence class $[v]$. If $[v]$ is abelian, we know by Proposition 4.4 of [CV08] that

$$Out(A_\Gamma) = Tr \rtimes (GL(A_{[v]}) \times Out(A_{lk[v]}))$$

where Tr is the free abelian group generated by the leaf transvections. Since $[v]$ is abelian, $GL(A_{[v]}) = GL(k, \mathbb{Z})$, which is residually finite, and $Out(A_{lk[v]})$ is residually finite by induction. The result now follows because semi-direct products of finitely generated residually finite groups are residually finite [Mi71]. If $[v]$ is non-abelian, then $Out(A_\Gamma) = Out(A_{[v]}) \times Out(A_{lk[v]})$ (or possibly a $\mathbb{Z}/2\mathbb{Z}$ -extension of this). Since $A_{[v]}$ is a free group, $Out(A_{[v]})$ is residually finite and $Out(A_{lk[v]})$ is residually finite by induction, so this case also follows.

Now suppose that Γ is not the star of a single equivalence class. Since $Out^0(A_\Gamma)$ has finite index in $Out(A_\Gamma)$, it suffices to prove that $Out^0(A_\Gamma)$ is residually finite. For any maximal equivalence class $[v]$, $Out^0(st[v])$ is residually finite by induction, so any element of $Out^0(A_\Gamma)$ which maps non-trivially under R is detectable by a finite group. It remains to show that the same is true for elements in the kernel K_R of R .

Let ϕ be an element of K_R . It follows from Lemma 3.3 and the fact that K_R is abelian that ϕ can be factored as

$$\phi = \phi_1 \circ \cdots \circ \phi_k$$

where ϕ_i is a product of \hat{v}_i -component conjugations, and the classes $[v_1], \dots, [v_k]$ are distinct. Let $[w]$ be a maximal vertex adjacent to $[v_1]$. Consider the image of ϕ under the

exclusion homomorphism $E_w : \text{Out}^0(A_\Gamma) \rightarrow \text{Out}^0(A_{\Gamma-[w]})$. By induction, the target group $\text{Out}^0(A_{\Gamma-[w]})$ is residually finite, so it suffices to show that this image, $\bar{\phi}$, is non-trivial.

Write $\bar{\phi} = \bar{\phi}_1 \circ \cdots \circ \bar{\phi}_k$. Note that $\bar{\phi}_1$ is still a nontrivial partial conjugation on $A_{\Gamma-[w]}$ since the vertices which were removed commuted with all elements of $[v_1]$. Moreover, for $i > 1$, the partial conjugations in $\bar{\phi}_i$ are either trivial, or are partial conjugations by elements distinct from $[v_1]$. It follows that $\bar{\phi}$ acts non-trivially on $A_{\Gamma-[w]}$ as required. \square

5. HOMOGENEOUS GRAPHS AND THE TITS ALTERNATIVE

Recall that the Tits alternative for a group G states that every subgroup of G is either virtually solvable or contains a non-abelian free group. Both $GL(n, \mathbb{Z})$ and $\text{Out}(F_n)$ are known to satisfy the Tits alternative [Ti72], [BFH00, BFH05]. We will show that $\text{Out}(A_\Gamma)$ satisfies the Tits alternative for a large class of graphs Γ .

Definition 5.1. Let Γ be a finite simplicial graph. We say Γ is *homogeneous of dimension 1* if it is non-empty and discrete (no edges). For $n > 1$, we say Γ is *homogeneous of dimension n* if it is connected and the link of every vertex is homogeneous of dimension $n - 1$.

It follows by a simple inductive argument that if Γ is homogeneous of dimension n then every maximal clique in G is an n -clique (hence the terminology ‘‘homogeneous’’). It also follows by induction that the link of any k -clique is a homogeneous graph of dimension $n - k$.

Examples 5.2. (1) For $n = 2$, a graph Γ is homogeneous if and only if Γ is connected and triangle-free. These are precisely the RAAGs studied in [CCV07]. (2) The join of two homogeneous graphs is again homogeneous so, for example, the join of two connected, triangle-free graphs is homogeneous of degree 4. (3) If Γ is the 1-skeleton of a connected triangulated n -manifold, then Γ is homogeneous of dimension n .

Our main concern is to be able to do inductive arguments on links of vertices; in particular, we will need such links to be connected or discrete at all stages of the induction. It may appear that homogeneity is a stronger condition than necessary. This is not the case.

Lemma 5.3. Γ is homogeneous of dimension $n > 1$ if and only if Γ is connected and the link of every (non-maximal) clique is either discrete or connected.

Proof. If Γ is homogeneous, then so is the link of every k -clique, $k < n$, so by definition it is either discrete or connected.

Conversely, assume that Γ is connected and the link of every non-maximal k -clique is either discrete or connected. We proceed by induction on the maximal size m of a clique in G . If $m = 2$, then the link of every vertex (1-clique) in Γ is discrete, so by definition, Γ is homogeneous of dimension 2.

For $m > 2$, we claim first that the link of every vertex is connected. For if Γ contains some vertex with a discrete link, then there exists an adjacent pair of vertices v, w such that the link of v is discrete while the link of w is not. In this case, v lies in $lk(w)$ but v is

not adjacent to any other vertex in $lk(w)$. This contradicts the assumption that the link of w is connected.

If Δ is a k -clique in $lk(v)$, then $\Delta * v$ is a $(k + 1)$ -clique in Γ . Since the link of Δ in $lk(v)$ is equal to the link of $\Delta * v$ in Γ , it is either discrete or connected. Thus, by induction, $lk(v)$ is homogeneous. Moreover, every link must be homogeneous of the same dimension, for if v, w are adjacent vertices, then the homogeneous dimension of $lk(v)$ and $lk(w)$ are both equal to $r - 1$ where r is the size of the maximal clique containing v and w . \square

The next lemma contains some other elementary facts about homogeneous graphs.

Lemma 5.4. *Let Γ be homogeneous of dimension n and assume that Γ is not the star of a single vertex (or equivalently, that Γ_0 contains more than one vertex). Let $[v]$ be a maximal equivalence class in Γ .*

- (1) *If $[v]$ is abelian, then $[v]$ is a singleton.*
- (2) *For any maximal $[v]$, $lk[v]$ is homogeneous of dimension $n - 1$ and is not the star of a single vertex.*

Proof. (1) Suppose $[v]$ is abelian and contains k vertices, so it spans a k -clique. Let Δ be an $n - k$ clique in $lk[v]$. Then $lk(\Delta)$ is homogeneous of dimension k and contains $[v]$. If $k > 1$, then $lk(\Delta)$ is connected, so either it consists of just the clique spanned by $[v]$, or it contains some other vertex w connected to $[v]$. In the former case, $\Gamma = [v] * lk[v] = st(v)$, which contradicts our hypothesis. In the latter case, w must be connected to every vertex in $[v]$ (since they are all equivalent), so $[v] \cup w$ spans a $k + 1$ -clique in $lk(\Delta)$ which is impossible. We conclude that $k = 1$, or in other words, $[v]$ is a singleton. This proves (1).

For (2), let $[v]$ be any maximal equivalence class. Then either $[v]$ is free, or a singleton and in either case, $lk[v] = lk(v)$, so it is homogeneous of dimension $n - 1$. If $lk[v]$ is contained in the star of a single vertex $w \in lk[v]$, then $[v] < [w]$. But this is impossible since $[v]$ is maximal. \square

We can now easily prove the main theorem of this section.

Theorem 5.5. *If Γ is homogeneous of dimension n , then $Out(A_\Gamma)$ satisfies the Tits alternative, that is, every subgroup of $Out(A_\Gamma)$ is either virtually solvable or contains a non-abelian free group.*

Proof. For Γ a complete graph, $Out(A_\Gamma) = GL(n, \mathbb{Z})$ so this follows from Tits' original theorem. So assume this is not the case. It suffices to prove the Tits Alternative for the finite index subgroup $Out^0(A_\Gamma)$. We proceed by induction on n . For $n = 1$, A_Γ is a free group and the theorem follows from [BFH00, BFH05].

If $n > 1$, then for every maximal $[v]$, $lk[v]$ is homogeneous of lower dimension, so by induction, $Out^0(A_{lk[v]})$ satisfies the Tits alternative. It is easy to verify that the Tits alternative is preserved under direct products, subgroups, and abelian extensions, so the theorem now follows from the exact sequence

$$1 \rightarrow K_P \rightarrow Out^0(A_\Gamma) \rightarrow \prod Out^0(A_{lk[v]}).$$

\square

6. SOLVABLE SUBGROUPS

6.1. Virtual derived length.

Definition 6.1. Let G be a solvable group and $G^{(i)}$ its derived series. The *derived length* of G is the least n such that $G^{(n)} = \{1\}$. The *virtual derived length* of G , which we denote by $vdl(G)$, is the minimum of the derived lengths of finite index subgroups of G .

For an arbitrary group H , define

$$\mu(H) = \max\{vdl(G) \mid G \text{ is a solvable subgroup of } H\}.$$

Note that if H is itself solvable, then $\mu(H) = vdl(H)$.

The following properties of $\mu(H)$ are easy exercises.

- Lemma 6.2.** (1) If $H = \prod H_i$, then $\mu(H) = \max\{\mu(H_i)\}$.
 (2) If N is a subgroup of H , then $\mu(N) \leq \mu(H)$. If $[H : N] < \infty$, then $\mu(N) = \mu(H)$.
 (3) If $N \triangleleft H$ is a solvable normal subgroup of derived length k , then $\mu(H) \leq \mu(H/N) + k$.

A group G has $vdl(G) = 1$ if and only if G is virtually abelian, and hence $\mu(H) = 1$ if and only if every solvable subgroup of H is virtually abelian. By [BFH05], $\mu(\text{Out}(F_n)) = 1$ for any free group F_n .

The situation for $GL(n, \mathbb{Z})$ is more complicated.

Proposition 6.3. Let U_n denote the subgroup of unitriangular matrices in $GL(n, \mathbb{Z})$. Then

$$\log_2(n) \leq \mu(U_n) \leq \mu(GL(n, \mathbb{Z})) < \log_2(n) + 2.$$

Proof. Let $e_{i,j}^a$ denote the elementary matrix with a in the (i, j) -th entry. For any finite index subgroup G of U_n , there exists $m \in \mathbb{Z}$ such that G contains all of the elementary matrices $e_{i,j}^m$ with $j > i$. The relation

$$[e_{i,j}^m, e_{k,j}^m] = e_{i,k}^{m^2}$$

then implies that the k th commutator subgroup $G^{(k)}$ contains all of the elementary matrices of the form $e_{i,j}^a$ with $j \geq i + 2^k$ and $a = m^{(2^k)}$. In particular, $G^{(k)}$ is non-trivial if $2^k < n$. The first inequality of the proposition follows.

The second inequality follows from Lemma 6.2(2). For the last inequality, we use a theorem of Mal'cev [Ma56], which implies that every solvable subgroup $H \subset GL(n, \mathbb{Z})$ is virtually isomorphic to a subgroup of $T_n(\mathcal{O})$, the upper triangular matrices over the ring of integers \mathcal{O} in some number field. The first commutator subgroup of $T_n(\mathcal{O})$ lies in $U_n(\mathcal{O})$ and it is easy to verify that $U_n(\mathcal{O})$ has derived length less than $\log_n(n) + 1$. It follows that

$$vdl(H) \leq dl(T_n(\mathcal{O})) < \log_2(n) + 2.$$

Since this holds for every solvable subgroup of $GL(n, \mathbb{Z})$, the last inequality of the proposition follows. \square

6.2. Maximum derived length for homogeneous graphs. In the case of a homogeneous graph Γ , it is easy to obtain an upper bound on the virtual derived length of solvable subgroups of $Out(A_\Gamma)$:

Theorem 6.4. *Suppose Γ is homogeneous of dimension n . Then $\mu(Out(A_\Gamma)) \leq n$.*

Proof. If Γ is a complete graph, $Out(A_\Gamma) \cong GL(n, \mathbb{Z})$, which has virtual derived length $\mu(GL(n, \mathbb{Z})) < \log_2(n) + 2 \leq n + 1$. So we may assume Γ is not a complete graph, in which case it contains a maximal vertex v with $lk[v]$ non-empty.

Since $Out^0(A_\Gamma)$ has finite index in $Out(A_\Gamma)$, their maximal derived lengths μ agree. We proceed by induction on n . For $n = 1$, Γ is discrete so $\mu(Out(A_\Gamma)) = 1$ by [BFH05]. For $n > 1$, we apply Lemma 6.2 to the abelian extension,

$$1 \rightarrow K_P \rightarrow Out^0(A_\Gamma) \rightarrow \prod Out^0(A_{lk[v]})$$

to conclude that $\mu(Out^0(A_\Gamma)) \leq 1 + \max\{\mu(Out^0(A_{lk[v]}))\}$. By Lemma 5.4, $lk[v]$ is homogeneous of dimension $n - 1$, so the theorem follows by induction. \square

Here is a stronger formulation of the previous theorem. If Γ is homogeneous of dimension n and Δ is an $(n - 1)$ -clique, then $lk(\Delta)$ is discrete, hence generates a free group $F(lk(\Delta))$.

Theorem 6.5. *Let Γ be homogeneous of dimension $n \geq 2$. Then there is a homomorphism*

$$Q : Out^0(A_\Gamma) \rightarrow \prod Out(F(lk(\Delta))),$$

where the product is taken over some collection of $(n - 1)$ -cliques, such that the kernel of Q is a solvable group of derived length at most $n - 1$.

Proof. Induction on n . For $n = 2$, take $Q = P$. The kernel K_P is abelian.

Suppose $n > 2$. Then P maps $Out^0(A_\Gamma)$ to a product of groups $Out^0(A_{lk[v]})$, where $lk[v]$ is homogeneous of dimension $n - 1$. By induction, there exists a homomorphism Q_v from $Out^0(A_{lk[v]})$ to a product of groups $Out(F(lk_v(\Delta)))$ where Δ is an $n - 2$ clique in $lk(v)$ and $lk_v(\Delta)$ is its link. The kernel H_v of Q_v is solvable of derived length at most $n - 2$.

Let $\Delta' = \Delta * v$. Then Δ' is an $(n - 1)$ -clique in Γ whose link $lk(\Delta')$ is exactly $lk_v(\Delta)$. Thus the composite $Q = (\prod Q_v) \circ P$ gives the desired homomorphism. The kernel of Q fits in an exact sequence

$$1 \rightarrow K_P \rightarrow \ker Q \rightarrow \prod H_v.$$

It follows that $\ker Q$ is solvable of derived length at most $n - 1$. \square

6.3. Examples of solvable subgroups. We now investigate lower bounds on the virtual derived length of $Out(A_\Gamma)$. If $[v]$ is an abelian equivalence class with k elements, then $GL(k, \mathbb{Z})$ embeds as a subgroup of $Out(A_\Gamma)$; in particular, $Out(A_\Gamma)$ contains solvable subgroups of virtual derived length at least $\log_2(k)$. In homogeneous graphs, abelian equivalence classes have only one element, so one cannot construct non-abelian solvable subgroups in this way. However, non-abelian solvable subgroups do exist, and we show two ways of constructing them in this section. More examples may be found in [Da09].

Let U_n denote the unitriangular matrices in $GL(n, \mathbb{Z})$, that is, the (lower) triangular matrices with 1's on the diagonal.

Proposition 6.6. *Let Γ be any finite simplicial graph. Suppose Γ contains $k + 1$ distinct vertices, v_0, \dots, v_k satisfying*

- (1) v_1, \dots, v_k span a k -clique
- (2) $[v_0] \leq [v_1] \leq \dots \leq [v_k]$

Then $Out(A_\Gamma)$ contains a subgroup isomorphic to U_{k+1} . In particular, $\mu(Out(A_\Gamma)) \geq \log_2(k + 1)$.

Proof. Let α_i denote the transvection $v_{i-1} \mapsto v_{i-1}v_i$. Let H denote the subgroup of $Out(A_\Gamma)$ generated by α_i , $1 \leq i \leq k$. Since H preserves the subgraph Γ' spanned by the v_i 's, it restricts to a subgroup of $Out(A_{\Gamma'})$. It is easy to see that this restriction maps H isomorphically onto its image, so WLOG, we may assume that $\Gamma = \Gamma'$.

Abelianizing A_Γ gives a map ρ from H to $GL(k + 1, \mathbb{Z})$. The image of α_i under ρ is the elementary matrix $e^{i+1, i}$. It follows that the image of H in $GL(k + 1, \mathbb{Z})$ is precisely U_{k+1} . Thus, it suffices to verify that the kernel of ρ is trivial. Let Δ be the clique spanned by v_1, \dots, v_k . Note that an element of H takes each v_i to $v_i w_i$ for some $w_i \in A_\Delta$. Since A_Δ is already abelian, a non-trivial w_i cannot be killed by abelianizing A_Γ . Thus, ρ is injective. \square

Proposition 6.7. *Let Γ be any finite simplicial graph. Suppose Γ contains k distinct vertices, v_1, \dots, v_k satisfying*

- (1) v_1, \dots, v_k span a k -clique
- (2) $[v_1] \leq \dots \leq [v_k]$
- (3) $\Gamma - st(v_1)$ has at least two distinct components that are not contained in $st(v_k)$

Then $Out(A_\Gamma)$ contains a subgroup isomorphic to U_{k+1} . In particular, $\mu(Out(A_\Gamma)) \geq \log_2(k + 1)$.

Proof. For $i = 2, \dots, k$, take α_i to be the transvection $v_{i-1} \mapsto v_{i-1}v_i$. Let C be a component of $\Gamma - st(v_1)$ which is not contained in $st(v_k)$. For $i = 1, \dots, k$, take β_i to be the partial conjugation of C by v_i . Note that β_i is non-trivial in $Out(A_\Gamma)$ since condition (3) guarantees that $\Gamma - st(v_i)$ contains at least two components. Let H denote the subgroup of $Out(A_\Gamma)$ generated by the α_i 's and β_i 's. We claim that H is isomorphic to U_{k+1} .

Since α_i acts only on vertices in $st(v_1)$ while β_i acts only on vertices not in $st(v_1)$, the subgroups H_α and H_β generated by the α_i 's and β_i 's respectively, are disjoint and H_β is easily seen to be normal in H . Hence H is the semi-direct product, $H = H_\beta \rtimes H_\alpha$. The subgroup H_α is isomorphic to U_k , as shown in the proof of the previous proposition, while H_β is isomorphic to the free abelian group A_Δ generated by the clique Δ spanned by the v_i 's. It is now easy to verify that H is isomorphic to U_{k+1} as claimed. \square

Example 6.8. Suppose A_Γ is homogeneous of dimension 2. Then by Theorem 6.4, $\mu(Out(A_\Gamma)) \leq 2$. In the next section (Corollary 6.13), we will show that if Γ has no leaves, then $\mu(Out(A_\Gamma)) = 1$. If Γ does have leaves, and for some leaf v , $st(v)$ separates Γ , then $\mu(Out(A_\Gamma)) = 2$. For if the components of $\Gamma - st(v)$ are not leaves, then Proposition 6.7 implies that $\log_2(3) \leq \mu(Out(A_\Gamma))$, and if some component is a leaf v' attached at the same base w , then Proposition 6.6 applied to $[v] \leq [v'] \leq [w]$ gives the same result.

6.4. Translation lengths and solvable subgroups. In constructing the non-abelian solvable subgroups above, a key role was played by transvections $v \rightarrow vw$ between adjacent (i.e. commuting) vertices. We call these *adjacent transvections*. In this section, we will show that without adjacent transvections, no such subgroups can exist for homogenous graphs.

Recall that $Out(A_\Gamma)$ is generated by the finite set S consisting of graph symmetries, inversions, partial conjugations and transvections. Define the following subsets of S ,

$$\begin{aligned}\tilde{S} &= S - \{\text{adjacent transvections}\} \\ \tilde{S}^0 &= \tilde{S} - \{\text{graph symmetries}\}\end{aligned}$$

and the subgroups of $Out(A_\Gamma)$ generated by them,

$$\begin{aligned}\tilde{Out}(A_\Gamma) &= \langle \tilde{S} \rangle \\ \tilde{Out}^0(A_\Gamma) &= \langle \tilde{S}^0 \rangle.\end{aligned}$$

We will prove that for Γ homogeneous, all solvable subgroups of $\tilde{Out}(A_\Gamma)$ are virtually abelian.

The proof proceeds by studying the translation lengths of infinite-order elements. The connection between solvable subgroups and translation lengths was first pointed out by Gromov [Gr87].

Definition 6.9. Let G be a group with finite generating set S , and let $\|g\|$ denote the word length of g in S . The *translation length* $\tau(g) = \tau_{G,S}(g)$ is the limit

$$\lim_{k \rightarrow \infty} \frac{\|g^k\|}{k}.$$

Elementary properties of translation lengths include the following (see [GS91], Lemma 6.2):

- $\tau(g^k) = k\tau(g)$.
- If S' is a different finite generating set, then $\tau_{G,S}(g)$ is positive if and only if $\tau_{G,S'}(g)$ is positive.
- If $H \leq G$ is a finitely-generated subgroup, and the generating set for G includes the generating set for H , then $\tau_H(h) \geq \tau_G(h)$.

Note that the kernel K_R of the amalgamated restriction homomorphism lies in $\tilde{Out}^0(A_\Gamma)$ since it is generated by products of partial conjugations.

Proposition 6.10. *Assume Γ is connected. Then every element of the kernel K_R of the amalgamated restriction homomorphism has positive translation length in $\tilde{Out}^0(A_\Gamma)$.*

Proof. Fix $\phi \in K_R$. We will find a function $\lambda = \lambda_\phi: \tilde{Out}^0(A_\Gamma) \rightarrow \mathbb{R}_{\geq 0}$ satisfying

- (1) $\lambda(\phi^k) \geq \frac{k}{2}$ and
- (2) $\lambda(\gamma_1 \dots \gamma_k) \leq 2k$ for $\gamma_i \in \tilde{S}^0$

The proof is then finished by the following argument. Let $m_k = \|\phi^k\|$, and write $\phi^k = \gamma_1 \dots \gamma_{m_k}$ with $\gamma_i \in \tilde{S}^0$. Then

$$\frac{k}{2} \leq \lambda(\phi^k) = \lambda(\gamma_1 \dots \gamma_{m_k}) \leq 2m_k$$

so

$$\frac{m_k}{k} \geq \frac{1}{4} > 0$$

for all k , so

$$\tau(\phi) = \lim_{k \rightarrow \infty} \frac{m_k}{k} \geq \frac{1}{4} > 0.$$

To define λ recall that by Theorem 3.3, the kernel K_R is free abelian, generated by \hat{w} -component conjugations. We write

$$\phi = \phi_{w_1} \phi_{w_2} \dots \phi_{w_k}$$

where ϕ_{w_i} is a nontrivial product of conjugations by w_i and the w_i are distinct.

First observe that the only transvections onto w_i are adjacent transvections. For if u is not adjacent to w_i and $w_i \leq u$, then there is only one non-trivial \hat{w}_i -component (the component of u), hence the unique \hat{w}_i -component conjugation is an inner automorphism. It follows that every element of $\tilde{Out}^0(A_\Gamma)$ fixes w_i up to conjugacy and inversion.

Set $w = w_1$. For an arbitrary element $x \in A_\Gamma$, define $p(x) = p_w(x)$ to be the absolute value of the largest power of w which can occur in a minimal-length word representing x . For example, if u or v does not commute with w , then $p(wuvw^{-2}) = 2$. If a minimal word representing x does not contain any powers of w , then $p(x) = 0$.

In [HM95], Hermiller and Meier describe a "left greedy" normal form for words in A_Γ , obtained by shuffling letters as far left as possible using the commuting relations and canceling inverse pairs whenever they occur. In particular, any reduced word can be put in normal form just by shuffling. It follows that the highest power of w that can occur in a minimal word for x is equal to the highest power of w appearing in the normal form for x .

For any automorphism $f \in Aut(A_\Gamma)$, define $p(f)$ to be the maximum over all vertices $v \neq w$ of $p(f(v))$. For an outer automorphism $\phi \in \tilde{Out}^0(A_\Gamma)$, define $\lambda(\phi)$ to be the minimum value of $p(f)$ as f ranges over automorphisms f representing ϕ . We must show that λ satisfies properties (1) and (2) above.

(1) Let f_w be a \hat{w} -component conjugation on the \hat{w} component C , and let $v \neq w$ be a vertex of Γ . If $v \in C - st(w)$, then $p(f_w^k(v)) = k$, and $p(f_w^k(v)) = 0$ otherwise. An inner automorphism can reduce the power of w by shifting it to vertices in the complement of C , but cannot reduce the maximum power of w over all vertices by more than $\lfloor k/2 \rfloor$. Since $\phi_w = \phi_{w_1}$ is non-trivial on at least one \hat{w} -component, this implies $\lambda(\phi_w^k) \geq k/2$. Since the partial conjugations ϕ_i for $i > 1$ do not change the power of w occurring at any vertex, we conclude that $\lambda(\phi^k) \geq k/2$.

(2) To prove property (2) we need to first establish some properties of the power function p . By abuse of notation, we will view \tilde{S}^0 as a subset of $Aut(A_\Gamma)$ in the obvious way.

Claim. Let $x \in A_\Gamma$. If $p(x) = 0$ and $f \in \tilde{S}^0$ then $p(f(x)) \leq 1$.

Proof. If f is a transvection or partial conjugation by some $u \neq w$, then $p(f(x)) = 0$. Likewise for inversions. So the only cases we have to consider are when f is either a non-adjacent transvection of w onto v or partial conjugation of $C \subset \Gamma$ by w .

Suppose f is a (non-adjacent) transvection $f: v \mapsto vw$ or $f: v \mapsto wv$. Then $f(x)$ has the property that any two copies of w are separated by v and any two copies of w^{-1} are separated by v^{-1} . ‘‘Shuffling left’’ can never switch the order of v and w , so this must also be true in the normal form for $f(x)$.

If f is a partial conjugation by w , then the w ’s in $f(x)$ alternate, i.e.

$$f(x) = a_1 w a_2 w^{-1} a_3 w \dots$$

where the a_i are words which do not use w or w^{-1} , so shuffling left can only cancel w -pairs, never increase the power to more than 1. \square

A minimal word representing $x \in A_\Gamma$ can be put in the form $a_0 w^{k_1} a_1 w^{k_2} \dots w^{k_n} a_n$ where

- a_i contains no w and
- w does not commute with a_i for $1 \leq i \leq n - 1$.

so that $p(x) = \max\{k_i\}$.

Claim. For any $f \in \tilde{S}^0$ and $x \in A_\Gamma$, $p(f(x)) \leq p(x) + 2$.

Proof. First assume that $f(w) = w$. This holds for all generators in \tilde{S}^0 with the exception of a partial conjugation by u of a component C containing w . Write $x = a_0 w^{k_1} a_1 w^{k_2} \dots w^{k_n} a_n$ as above. Let b_i be the normal form for $f(a_i)$. Then $f(x) = b_0 w^{k_1} b_1 w^{k_2} \dots w^{k_n} b_n$, where b_i does not commute with w .

Case 1: f is a partial conjugation or transvection by w . Then no w can shuffle across an entire b_i , so we need only consider the highest power appearing in $b_{i-1} w^{k_i} b_i$. Now by the previous claim, b_i is of the form

$$b_i = c_1 w^{\pm 1} c_2 w^{\pm 1} \dots w^{\pm 1} c_k$$

where w does not commute with c_2, \dots, c_{k-1} and similarly for b_{i-1} . It follows that left shuffling of $b_{i-1} w^{k_i} b_i$ can at worst combine w^{k_i} with the last w in b_{i-1} and the first w in b_i , producing a power of at most $|k_i| + 2$.

Case 2: f is a partial conjugation (on a component not containing w) or transvection by some $u \neq w$. Then no new w ’s appear and no w can shuffle across an entire b_i , so the maximum power of w does not change, i.e., $p(f(x)) = p(x)$.

It remains to consider the case where f is a partial conjugation by u with $f(w) = u w u^{-1}$. Then f can be written as the composite of an inner automorphism by u followed by a product of partial conjugations fixing w . By case 2 above, we have $p(f(x)) = p(uxu^{-1})$. Since u does not commute with w , conjugating by u changes only the factors a_0 and a_n in the normal form for x . Thus $p(uxu^{-1}) = p(x)$. \square

If $f \in \text{Aut}(A_\Gamma)$ can be written as a product of m elements of \tilde{S}^0 , then the above claim shows that $p(f(v)) \leq 2m$, for any vertex $v \neq w$. If $\phi \in \tilde{\text{Out}}^0(A_\Gamma)$ can be written as a product $\phi = \phi_1 \dots \phi_m$, with ϕ_i represented by $f_i \in \tilde{S}^0$, then $\lambda(\phi) \leq p(f) \leq 2m$. This completes the proof of the proposition. \square

Corollary 6.11. *If Γ is homogeneous of dimension n , then the translation length of every infinite-order element of $\tilde{Out}^0(A_\Gamma)$ is positive.*

Proof. We proceed by induction on n . For $n = 1$, A_Γ is a free group F_k and $\tilde{Out}^0(A_\Gamma) = Out(F_k)$. Alibegovic proved that infinite order elements of $Out(F_k)$ have positive translation length [Al02].

For $n > 1$, we will make use of the projection homomorphism

$$P = \prod P_v: Out^0(A_\Gamma) \rightarrow \prod_{v \text{ maximal}} Out(A_{lk[v]}).$$

Note that each projection P_v maps generators in \tilde{S}^0 to either the trivial map or to a generator of the same form in $Out(A_{lk[v]})$. Thus the image of $\tilde{Out}^0(A_\Gamma)$ lies in the product of the subgroups $\tilde{Out}^0(A_{lk[v]})$. Moreover, the kernel of P restricted to $\tilde{Out}^0(A_\Gamma)$ is just K_R . This follows from the fact that K_P is generated by K_R and leaf-transvections, which by definition, are adjacent transvections.

By [CV08], all of the groups we are considering are virtually torsion-free. Let $G \leq \tilde{Out}^0(A_\Gamma)$ be the inverse image of a torsion-free finite-index subgroup of $\prod \tilde{Out}^0(A_{lk[v]})$. If $\phi \in \tilde{Out}^0(A_\Gamma)$ has infinite order, then some power of ϕ is a non-trivial (infinite-order) element in G , so we need only prove that elements of G have positive translation length in $\tilde{Out}^0(A_\Gamma)$.

If the image of $\phi \in G$ is non-trivial in some $\tilde{Out}^0(A_{lk[v]})$, then it has positive translation length by induction. If the image is trivial, then ϕ lies in K_R so we are done by Proposition 6.10. \square

Corollary 6.12. *If Γ is homogeneous of dimension n , then $\tilde{Out}(A_\Gamma)$ satisfies the strong Tits alternative, that is, every subgroup of $\tilde{Out}(A_\Gamma)$ is either virtually abelian or contains a non-abelian free group.*

Proof. By Theorem 5.5, every subgroup not containing a free group is virtually solvable. So it remains to show that every solvable subgroup is virtually abelian. Since $\tilde{Out}^0(A_\Gamma)$ has finite index in $\tilde{Out}(A_\Gamma)$, it suffices to prove the same statement for $\tilde{Out}^0(A_\Gamma)$.

Bestvina [Be99], citing arguments from Conner [Co00] and Gersten and Short [GS91], shows that if a finitely-generated group is positive, virtually torsion-free and its abelian subgroups are finitely generated, then solvable subgroups must be virtually abelian. $\tilde{Out}^0(A_\Gamma)$ is positive by Corollary 6.11 and virtually torsion-free by [CV08]. Since Γ is homogeneous, the fact that abelian subgroups of $Out^0(A_\Gamma)$ are finitely generated follows by a simple induction from the same fact for $Out(F_n)$ and $GL(n, \mathbb{Z})$ using the projection homomorphisms. \square

In dimension 2, the only adjacent transvections are leaf transvections, so if Γ has no leaves, then $\tilde{Out}(A_\Gamma) = Out(A_\Gamma)$. Thus the following is a special case of Corollary 6.12.

Corollary 6.13. *If Γ is connected with no triangles and no leaves, then $Out(A_\Gamma)$ satisfies the strong Tits alternative.*

7. QUESTIONS

Since the projection homomorphism $P: \text{Out}^0(A_\Gamma) \rightarrow \prod \text{Out}^0(A_{lk[v]})$ is defined only for connected graphs Γ , inductive arguments using P break down if the links of maximal vertices are not connected, unless the desired result is known by some other argument for outer automorphism groups of free products. For homogeneous graphs, the links are always connected so this is not an issue, but several of the questions answered in this paper remain open for non-homogeneous graphs. Specifically, we can ask

- (1) Is the maximal virtual derived length of a solvable subgroup of $\text{Out}(A_\Gamma)$ bounded by the dimension of A_Γ ?
- (2) Does $\text{Out}(A_\Gamma)$ satisfy the Tits alternative?

REFERENCES

- [Al02] Emina Alibegovich, *Translation lengths in $\text{Out}(F_n)$* , *Geom. Dedicata* **92** (2002) 87–93
- [Be99] Mladen Bestvina, *Non-positively curved aspects of Artin groups of finite type*, *Geom. Topol.* **3** (1999) 269–302
- [BFH00] Mladen Bestvina, Mark Feighn and Michael Handel, *The Tits alternative for $\text{Out}(F_n)$ I: Dynamics of exponentially-growing automorphisms*, *Annals of Mathematics* **151** (2000), 517–623.
- [BFH05] Mladen Bestvina, Mark Feighn and Michael Handel, *The Tits alternative for $\text{Out}(F_n)$. II. A Kolchin type theorem*, *Ann. of Math. (2)* **161** (2005) 1–59
- [BCV09] Kai-Uwe Bux, Ruth Charney and Karen Vogtmann, *Automorphisms of two-dimensional RAAGs and partially symmetric automorphisms of free groups*, to appear in *Groups, dynamics and*
- [Ch07] Ruth Charney, *An introduction to right-angled Artin groups*, *Geom. Dedicata* **125** (2007) 141–158
- [CCV07] Ruth Charney, John Crisp and Karen Vogtmann, *Automorphisms of 2-dimensional right-angled Artin groups*, *Geom. and Topology* **11** (2007), 2227–2264.
- [CV08] Ruth Charney and Karen Vogtmann, *Finiteness properties of automorphism groups of right-angled Artin groups*, *Bulletin LMS* (2009); doi: 10.1112/blms/bdn108
- [Co00] Greg Connor, *Discreteness properties of translation numbers in solvable groups*, *J. Group Theory* **3** (2000), no. 1, 77–94.
- [Da09] Matt Day, *On solvable automorphism groups of RAAGs*, preprint.
- [GS91] Steve Gersten and Hamish Short, *Rational subgroups of biautomatic groups*, *Ann. of Math. (2)* **134** (1) (1991), 125–158
- [Gr74] Edna Grossman, *On the residual finiteness of certain mapping class groups*, *J. London Math. Soc. (2)*, **9** (1974), 160–164
- [Gr87] M. Gromov, *Hyperbolic groups*, in *Essays on Group Theory*, MSRI series, vol , edited by S. Gersten, Springer-Verlag, 1987.
- [GL07] Vincent Guirardel and Gilbert Levitt, *The Outer space of a free product*, *Proc. Lond. Math. Soc. (3)* **94** (2007), 695–714
- [HM95] S. Hermiller and J. Meier, *Algorithms and geometry for graph products of groups*, *J. Algebra* **171** (1995), no. 1, 230257
- [Lau95] MR Laurence, *A generating set for the automorphism group of a graph group*, *J. London Math. Soc. (2)* **52** (1995) 318–334
- [Ma56] A. I. Mal'cev, *On certain classes of innite soluble groups*, *Mat. Sbornik* **28** (1951) 567–588 (Russian); *Amer. Math. Soc. Translations (2)* **2** (1956) 1–21.
- [Mi71] Charles F. Miller, *On Group-Theoretic Decision Problems and their Classification*, Princeton Univ. Press, Princeton, NJ, 1971.
- [Mi09] A. Minasyan, *Hereditary conjugacy separability of right angled Artin groups and its applications*, arXiv:0905.1282

- [MiOs09] A. Minasyan, D. Osin, *Normal automorphisms of relatively hyperbolic groups*, arXiv:0809.2408
- [Ser89] H Servatius, *Automorphisms of graph groups*, J. Algebra 126 (1989) 34–60
- [Ti72] Jacques Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250270