

# A NOTE ON 2-DISTANT NONCROSSING PARTITIONS AND WEIGHTED MOTZKIN PATHS

IRA M. GESSEL AND JANG SOO KIM

ABSTRACT. We prove a conjecture of Drake and Kim: the number of 2-distant noncrossing partitions of  $\{1, 2, \dots, n\}$  is equal to the sum of weights of Motzkin paths of length  $n$ , where the weight of a Motzkin path is a product of certain fractions involving Fibonacci numbers. We provide two proofs of their conjecture: one uses continued fractions and the other is combinatorial.

## 1. INTRODUCTION

A *Motzkin path* of length  $n$  is a lattice path from  $(0, 0)$  to  $(n, 0)$  consisting of up steps  $U = (1, 1)$ , down steps  $D = (1, -1)$  and horizontal steps  $H = (1, 0)$  that never goes below the  $x$ -axis. The *height* of a step in a Motzkin path is the  $y$  coordinate of the ending point.

Given two sequences  $b = (b_0, b_1, \dots)$  and  $\lambda = (\lambda_0, \lambda_1, \dots)$ , the *weight* of a Motzkin path with respect to  $(b, \lambda)$  is the product of  $b_i$  and  $\lambda_i$  for each horizontal step and down step of height  $i$  respectively, see Figure 1. Let  $\text{Mot}_n(b, \lambda)$  denote the sum of weights of Motzkin paths of length  $n$  with respect to  $(b, \lambda)$ . This sum is closely related to orthogonal polynomials; see [5, 6].

Drake and Kim [1] defined the set  $\text{NC}_k(n)$  of  $k$ -distant noncrossing partitions of  $[n] = \{1, 2, \dots, n\}$ . For  $k \geq 0$ , a  $k$ -distant noncrossing partition is a set partition of  $[n]$  without two arcs  $(a, c)$  and  $(b, d)$  satisfying  $a < b \leq c < d$  and  $c - b \geq k$ , where an *arc* is a pair  $(i, j)$  of integers contained in the same block which does not contain any integer between them. For example,  $\pi = \{\{1, 5, 7\}, \{2, 3, 6\}, \{4\}\}$  is a 3-distant noncrossing partition but not a 2-distant noncrossing partition because  $\pi$  has two arcs  $(1, 5)$  and  $(3, 6)$  with  $5 - 3 \geq 2$ . Note that the 1-distant noncrossing partitions are the ordinary noncrossing partitions, which implies that  $\#\text{NC}_1(n)$  is equal to the Catalan number  $\frac{1}{n+1} \binom{2n}{n}$ . It is not difficult to see that  $\text{NC}_0(n)$  is in bijection

---

The first author was supported by NSA Grant H98230-10-1-0196. The second author was supported by the grant ANR08-JCJC-0011.




FIGURE 1. A Motzkin path and the weights of its steps with respect to  $(b, \lambda)$ .

with the set of Motzkin paths of length  $n$ . In the same paper, they proved that

$$(1) \quad \sum_{n \geq 0} \# \text{NC}_2(n) x^n = \frac{3}{2} - \frac{1}{2} \sqrt{\frac{1-5x}{1-x}}.$$

The number  $\# \text{NC}_2(n)$  also counts many combinatorial objects: Schröder paths with no peaks at even levels, etc; see [2, 4, 7].

There are simple expressions of  $\# \text{NC}_k(n)$  using Motzkin paths for  $k = 0, 1, 3$ :

$$\begin{aligned} \# \text{NC}_0(n) &= \text{Mot}_n((1, 1, \dots), (1, 1, \dots)), \\ \# \text{NC}_1(n) &= \text{Mot}_n((1, 2, 2, \dots), (1, 1, \dots)), \\ \# \text{NC}_3(n) &= \text{Mot}_n((1, 2, 3, 3, \dots), (1, 2, 2, \dots)), \end{aligned}$$

where the second equation is well known and the third one was first conjectured by Drake and Kim [1] and proved by Kim [3]. The main purpose of this paper is to prove the following theorem which was also conjectured by Drake and Kim [1].

**Theorem 1.1.** *Let  $b = (b_0, b_1, \dots)$  and  $\lambda = (\lambda_0, \lambda_1, \dots)$  be the sequences with  $b_0 = \lambda_0 = 1$  and for  $n \geq 1$ ,*

$$(2) \quad b_n = 3 - \frac{1}{F_{2n-1}F_{2n-3}} \quad \text{and} \quad \lambda_n = 1 + \frac{1}{F_{2n-1}^2},$$

where  $F_m$  is the Fibonacci number defined by  $F_0 = 0, F_1 = 1$ , and  $F_m = F_{m-1} + F_{m-2}$  for all  $m$  (so  $F_{-1} = 1$ ). Then we have

$$\# \text{NC}_2(n) = \text{Mot}_n(b, \lambda).$$

Theorem 1.1 is very interesting because it is not even obvious that  $\text{Mot}_n(b, \lambda)$  is an integer. In this paper, we give two proofs of Theorem 1.1: one uses continued fractions and the other is combinatorial.

## 2. CONTINUED FRACTIONS

Let  $\alpha = (\alpha_0, \alpha_1, \alpha_2, \dots)$ ,  $\beta = (\beta_0, \beta_1, \beta_2, \dots)$ , and  $c = (c_0, c_1, c_2, \dots)$  be sequences of numbers.

Let  $J(x; \alpha_0, \beta_0; \alpha_1, \beta_1; \alpha_2, \beta_2; \dots) = J(x; \alpha, \beta)$  denote the *J-fraction*

$$\frac{1}{1 - \alpha_0 x - \frac{\beta_0 x^2}{1 - \alpha_1 x - \frac{\beta_1 x^2}{1 - \alpha_2 x - \dots}}}$$

and let  $S(x; c_0, c_1, c_2, \dots) = S(x; c)$  denote the *S-fraction*

$$\frac{1}{1 - \frac{c_0 x}{1 - \frac{c_1 x}{1 - \dots}}}$$

A *Dyck path* of length  $2n$  is a lattice path from  $(0, 0)$  to  $(2n, 0)$  consisting of up steps  $U = (1, 1)$  and down steps  $D = (1, -1)$  that never goes below the  $x$ -axis. The *height* of a step in a Dyck path is the  $y$  coordinate of the ending point. The *weight* of a Dyck path with respect to  $c$  is the product of  $c_i$  for each down step of height  $i$ , see Figure 2. Let  $\text{Dyck}_n(c)$  denote the sum of weights of Dyck paths of length  $2n$  with respect to  $c$ .




FIGURE 2. A Dyck path and the weights of its steps with respect to  $c$ .

It is well known that

$$\sum_{n \geq 0} \text{Mot}_n(\alpha, \beta)x^n = J(x; \alpha, \beta) \quad \text{and} \quad \sum_{n \geq 0} \text{Dyck}_n(c)x^n = S(x; c).$$

The following proposition is easy to see.

**Proposition 2.1.** *If  $\alpha_n = c_{2n-1} + c_{2n}$  and  $\beta_n = c_{2n}c_{2n+1}$  for all  $n \geq 0$ , with  $c_{-1} = 0$ , then  $S(x; c) = J(x; \alpha, \beta)$ .*

One can prove Proposition 2.1 by the following observation: a Motzkin path may be obtained from a Dyck path by taking steps two at a time and changing  $UU$ ,  $UD$ ,  $DU$  and  $DD$ , respectively, to  $U$ ,  $H$ ,  $H$  and  $D$ . For example, the Motzkin path in Figure 1 is obtained from the Dyck path in Figure 2 in this way.

Let  $d = (d_0, d_1, d_2, \dots)$  be the sequence with  $d_0 = 1$  and for  $n \geq 1$ ,

$$(3) \quad d_{2n-1} = \frac{F_{2n-1}}{F_{2n-3}}, \quad d_{2n} = \frac{1}{d_{2n-1}}.$$

Recall the two sequences  $b = (b_0, b_1, \dots)$  and  $\lambda = (\lambda_0, \lambda_1, \dots)$  defined in (2).

**Lemma 2.2.** *We have the following.*

- (1)  $b_n = d_{2n-1} + d_{2n}$  for all  $n \geq 0$ , where  $d_{-1} = 0$ .
- (2)  $\lambda_n = d_{2n}d_{2n+1}$  for all  $n \geq 0$ .
- (3)  $1/d_{2n-1} + d_{2n+1} = 3$  for all  $n \geq 1$ .

*Proof.* We will use two cases of the well-known Catalan identity for Fibonacci numbers,  $F_m^2 - F_{m+i}F_{m-i} = (-1)^{m-i}F_i^2$ .

(1) This is true for  $n = 0$ . For  $n \geq 1$  we have

$$\begin{aligned} d_{2n-1} + d_{2n} &= \frac{F_{2n-1}}{F_{2n-3}} + \frac{F_{2n-3}}{F_{2n-1}} = \frac{F_{2n-1}^2 + F_{2n-3}^2}{F_{2n-1}F_{2n-3}} = \frac{2F_{2n-1}F_{2n-3} + (F_{2n-1} - F_{2n-3})^2}{F_{2n-1}F_{2n-3}} \\ &= 2 + \frac{F_{2n-2}^2}{F_{2n-1}F_{2n-3}} = 3 + \frac{F_{2n-2}^2 - F_{2n-1}F_{2n-3}}{F_{2n-1}F_{2n-3}} = 3 - \frac{1}{F_{2n-1}F_{2n-3}} = b_n. \end{aligned}$$

(2) This is true for  $n = 0$ . For  $n \geq 1$  we have

$$d_{2n}d_{2n+1} = \frac{F_{2n-3}}{F_{2n-1}} \frac{F_{2n+1}}{F_{2n-1}} = \frac{F_{2n-1}^2 + (F_{2n-3}F_{2n+1} - F_{2n-1}^2)}{F_{2n-1}^2} = 1 + \frac{1}{F_{2n-1}^2} = \lambda_n.$$

(3) We have

$$\begin{aligned} \frac{1}{d_{2n-1}} + d_{2n+1} &= \frac{F_{2n-3}}{F_{2n-1}} + \frac{F_{2n+1}}{F_{2n-1}} = \frac{(F_{2n-1} - F_{2n-2}) + (F_{2n} + F_{2n-1})}{F_{2n-1}} \\ &= 2 + \frac{F_{2n} - F_{2n-2}}{F_{2n-1}} = 3. \end{aligned}$$

□

By Proposition 2.1 and Lemma 2.2, we obtain the following.

**Corollary 2.3.** *For the sequences  $b$ ,  $\lambda$  and  $d$  defined in (2) and (3), we have*

$$\text{Dyck}_n(d) = \text{Mot}_n(b, \lambda).$$

Now we can prove the following  $S$ -fraction formula for the generating function (1) for  $\#\text{NC}_2(n)$ .

**Theorem 2.4.** *We have*

$$\begin{aligned} \frac{3}{2} - \frac{1}{2} \sqrt{\frac{1-5x}{1-x}} &= S(x; 1, 1, 1, 2, \frac{1}{2}, \frac{5}{2}, \frac{2}{5}, \frac{13}{5}, \frac{5}{13}, \frac{34}{13}, \frac{13}{34}, \frac{89}{34}, \frac{34}{89}, \frac{233}{89}, \frac{89}{233}, \frac{610}{233}, \frac{233}{610}, \frac{1597}{610}, \dots) \\ &= S(x; d_0, d_1, d_2, \dots). \end{aligned}$$

To prove Theorem 2.4, we define  $R_n$  for  $n \geq -1$  by

$$\begin{aligned} R_{-1} &= \frac{3}{2} - \frac{1}{2} \sqrt{\frac{1-5x}{1-x}}, \\ R_{2n+1} &= d_{2n+1} + \frac{1-3x - \sqrt{(1-x)(1-5x)}}{2x}, \quad n \geq 0, \\ R_{2n} &= \frac{d_{2n}}{1-xR_{2n+1}}, \quad n \geq 0. \end{aligned}$$

One can easily check that  $R_m$  is a power series in  $x$  with constant term  $d_m$  (with  $d_{-1} = 1$ ), though this will follow from Lemma 2.5.

**Lemma 2.5.** *For  $m \geq -1$ , we have*

$$R_m = \frac{d_m}{1-xR_{m+1}},$$

where  $d_{-1} = 1$ .

*Proof.* By definition, this is true if  $m$  is even. Thus it is enough to prove that for  $n \geq 0$ ,

$$R_{2n-1} = \frac{d_{2n-1}}{1-xR_{2n}} = \frac{d_{2n-1}}{1 - \frac{xd_{2n}}{1-xR_{2n+1}}},$$

which is equivalent to

$$(4) \quad R_{2n+1} = \frac{1}{x} - \frac{1}{d_{2n-1}} - \frac{1}{R_{2n-1} - d_{2n-1}}.$$

We can check (4) directly for  $n = 0$ . Assume  $n \geq 1$ . Then the right-hand side of (4) is equal to

$$\begin{aligned} \frac{1}{x} - \frac{1}{d_{2n-1}} - \frac{2x}{1-3x - \sqrt{(1-x)(1-5x)}} &= \frac{1}{x} - \frac{1}{d_{2n-1}} - \frac{2x \left(1-3x + \sqrt{(1-x)(1-5x)}\right)}{(1-3x)^2 - (1-6x+5x^2)} \\ &= \frac{1}{x} - \frac{1}{d_{2n-1}} - \frac{1-3x + \sqrt{(1-x)(1-5x)}}{2x} \\ &= 3 - \frac{1}{d_{2n-1}} + \frac{1-3x - \sqrt{(1-x)(1-5x)}}{2x}. \end{aligned}$$

Since  $3 - 1/d_{2n-1} = d_{2n+1}$  by Lemma 2.2, we are done.  $\square$

*Proof of Theorem 2.4.* It follows from Lemma 2.5 that

$$\frac{3}{2} - \frac{1}{2} \sqrt{\frac{1-5x}{1-x}} = R_{-1} = \frac{1}{1-xR_0} = \frac{1}{1-\frac{d_0x}{1-xR_1}} = \frac{1}{1-\frac{d_0x}{1-\frac{d_1x}{1-xR_2}}} = \dots$$

Continuing, and taking a limit, gives the S-fraction for  $R_{-1}$ . □

By (1), Theorem 2.4 and Corollary 2.3, we obtain the following which proves Theorem 1.1.

$$\sum_{n \geq 0} \# \text{NC}_2(n)x^n = \frac{3}{2} - \frac{1}{2} \sqrt{\frac{1-5x}{1-x}} = \sum_{n \geq 0} \text{Dyck}_n(d)x^n = \sum_{n \geq 0} \text{Mot}_n(b, \lambda)x^n$$

### 3. A COMBINATORIAL PROOF

Let  $b$ ,  $\lambda$  and  $d$  be the sequences defined in (2) and (3).

Recall that in the previous section we have shown that  $\text{Dyck}_n(d) = \text{Mot}_n(b, \lambda)$  by changing a Dyck path of length  $2n$  to a Motzkin path of length  $n$ . We can do the same thing after deleting the first and the last steps of a Dyck path. More precisely, for a Dyck path of length  $2n$ , we delete the first and the last steps, take two steps at a time in the remaining  $2n - 2$  steps, and change  $UU$ ,  $UD$ ,  $DU$  and  $DD$ , respectively, to  $U$ ,  $H$ ,  $H$  and  $D$ . Then we obtain a Motzkin path of length  $n - 1$ . This argument shows that

$$(5) \quad \text{Dyck}_n(d) = d_0 \cdot \text{Mot}_{n-1}(\alpha, \beta) = \text{Mot}_{n-1}(\alpha, \beta),$$

where  $\alpha_n = d_{2n} + d_{2n+1}$  and  $\beta_n = d_{2n+1}d_{2n+2}$ . By (3) and Lemma 2.2, we have  $\alpha = (2, 3, 3, \dots)$  and  $\beta = (1, 1, \dots)$ . Note that we can also prove Theorem 2.4 using (5).

To find a connection between  $\text{Mot}_{n-1}(\alpha, \beta)$  and  $\text{NC}_2(n)$  we need the following definition.

A *Schröder path* of length  $2n$  is a lattice path from  $(0, 0)$  to  $(2n, 0)$  consisting of up steps  $U = (1, 1)$ , down steps  $D = (1, -1)$  and double horizontal steps  $H^2 = HH = (2, 0)$  that never goes below the  $x$ -axis. Let  $\text{SCH}_{\text{even}}(n)$  denote the set of Schröder paths of length  $2n$  such that all horizontal steps have even height.

**Proposition 3.1.** *Let  $\alpha = (2, 3, 3, \dots)$  and  $\beta = (1, 1, \dots)$ . Then, for  $n \geq 1$ , we have*

$$\text{Mot}_n(\alpha, \beta) = \# \text{SCH}_{\text{even}}(n).$$

*Proof.* From a Motzkin path of length  $n$  we obtain a Schröder path of length  $2n$  as follows. Change  $U$  and  $D$  to  $UU$  and  $DD$  respectively. For a horizontal step  $H$ , if its height is 0, we change it to either  $UD$  or  $HH$ , and if its height is greater than 0, we change it to either  $UD$ ,  $DU$  or  $HH$ . Then we get an element of  $\text{SCH}_{\text{even}}(n)$ . Since the weight of a horizontal step  $H$  in the Motzkin path is equal to the number of choices, the theorem follows. □

*Remark 1.* The definition of  $\text{SCH}_{\text{even}}(n)$  in [2] is the set of Schröder paths of length  $2n$  which have no peaks at even height. From such a path, by changing all the horizontal steps at odd height to peaks, we get a Schröder path whose horizontal steps are all at even height, and this transformation is easily seen to be a bijection.

Kim [2] found a bijection between  $\text{NC}_2(n)$  and  $\text{SCH}_{\text{even}}(n-1)$ . Using Kim's bijection in [2], Proposition 3.1, (5) and Corollary 2.3 we finally get the following sequence of identities which implies Theorem 1.1:

$$\#\text{NC}_2(n) = \#\text{SCH}_{\text{even}}(n-1) = \text{Mot}_{n-1}(\alpha, \beta) = \text{Dyck}_n(d) = \text{Mot}_n(b, \lambda).$$

#### ACKNOWLEDGEMENT

The authors would like to thank the anonymous referees for their careful reading and helpful comments which significantly improved the presentation of the paper.

#### REFERENCES

- [1] Dan Drake and Jang Soo Kim.  $k$ -distant crossings and nestings of matchings and partitions. *DMTCS Proceedings*, AK:349–360, 2009.
- [2] Jang Soo Kim. Bijections on two variations of noncrossing partitions. <http://arxiv.org/abs/0812.4091>.
- [3] Jang Soo Kim. Front representation of set partitions. <http://arxiv.org/abs/0907.1485>.
- [4] Toufik Mansour and Simone Severini. Enumeration of  $(k, 2)$ -noncrossing partitions. *Discrete Mathematics*, 308(20):4570–4577, October 2008.
- [5] Gérard Viennot. Une théorie combinatoire des polynômes orthogonaux généraux, Sept.–Oct. 1983. Notes from a conference given at the Université du Québec à Montréal.
- [6] Gérard Viennot. A combinatorial theory for general orthogonal polynomials with extensions and applications. In *Orthogonal polynomials and applications (Bar-le-Duc, 1984)*, volume 1171 of *Lecture Notes in Math.*, pages 139–157, Berlin, 1985. Springer.
- [7] Sherry Yan. Schröder paths and pattern avoiding partitions. *Int. Journal of Contemp. Math. Sciences*, Vol. 4, no. 17-20:979–986, 2009.

*E-mail address*, Ira M. Gessel: [gessel@brandeis.edu](mailto:gessel@brandeis.edu)

*E-mail address*, Jang Soo Kim: [kimjs@math.umn.edu](mailto:kimjs@math.umn.edu)