

# Solutions to review problems for Midterm 2


March 29, 2005

## 1 Solutions to problems from the book


**Problem 2.**  $D = \{(x, y, z) \mid z \geq x^2 + y^2\}$ , which is the set of points above the paraboloid  $z = x^2 + y^2$ .

**Problem 4.**  $z = \sqrt{x^2 + y^2 - 1}$ , so  $z \geq 0$  and  $x^2 + y^2 - z^2 = 1$ .

So the graph is the upper half of a hyperboloid of one sheet.



**Problem 8.**



**Problem 10.** as  $(x, y) \rightarrow (0, 0)$  Along the line  $x = y$ , the value of  $\frac{2xy}{x^2+2y^2}$  is  $\frac{2x^2}{3x^2}$ , so the limit as  $(x, y) \rightarrow (0, 0)$  along this line is  $\frac{2}{3}$ . In contrast, along the line  $x = -y$ ,  $\frac{2xy}{x^2+2y^2}$  is  $\frac{-2x^2}{3x^2} = -\frac{2}{3}$ , so the limit does not exist.

**Problem 16.**  $\frac{\partial w}{\partial x} = \frac{1}{y-z}$ ,  $\frac{\partial w}{\partial y} = \frac{-x}{(y-z)^2}$ , and  $\frac{\partial w}{\partial z} = \frac{x}{(y-z)^2}$ .

**Problem 22.**  $v_{rr} = 0$ ,  $v_{rs} = -\sin(s + 2t)$ ,  $v_{rt} = -2\sin(s + 2t)$ ,  $v_{ss} = -r \cos(s + 2t)$ ,  $v_{st} = -2r \cos(s + 2t)$ , and  $v_{tt} = -4r \cos(s + 2t)$ . The others are determined by these, by Clairaut's theorem.

**Problem 24.**  $\frac{\partial^2 \rho}{\partial x^2} = \frac{\partial}{\partial x}(x(x^2 + y^2 + z^2)^{-1/2}) = (x^2 + y^2 + z^2)^{-1/2} - x^2(x^2 + y^2 + z^2)^{-3/2} = \frac{x^2 + y^2 + z^2 - x^2}{(x^2 + y^2 + z^2)^{3/2}} = \frac{y^2 + z^2}{(x^2 + y^2 + z^2)^{3/2}}$ . The  $y$  and  $z$  derivatives are similar; adding them up together gives

$$\frac{(y^2 + z^2) + (x^2 + z^2) + (x^2 + y^2)}{(x^2 + y^2 + z^2)^{3/2}} = \frac{2(x^2 + y^2 + z^2)}{(x^2 + y^2 + z^2)^{3/2}} = \frac{2}{\rho}.$$

**Problem 28.** Write  $F(x, y, z) = xy + yz + zx - 3$ , so  $F_x = y + z$ ,  $F_y = x + z$ , and  $F_z = x + y$ . Thus the normal vector to the plane is  $\langle 2, 2, 2 \rangle$ , and the equation of the plane is  $2(x-1) + 2(y-1) + 2(z-1) = 0$ .

**Problem 36.**  $\frac{\partial z}{\partial x} = -y \sin xy - y \sin x$ , and  $\frac{\partial z}{\partial y} = -x \sin xy + \cos x$ . So

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial u} = (-y \sin xy - y \sin x)2u + (-x \sin xy + \cos x)(1).$$

Similarly,

$$\frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v} = (-y \sin xy - y \sin x)(1) + (-x \sin xy + \cos x)(-2v).$$

**Problem 42.** Use implicit differentiation. For the first, take the  $x$ -derivative of both sides:  $4yz^3 \frac{\partial z}{\partial x} + 2xz^3 + 3x^2 z^2 \frac{\partial z}{\partial x} = yze^{xyz} + xy \frac{\partial z}{\partial x} e^{xyz}$ . Collecting terms involving  $\frac{\partial z}{\partial x}$  gives

$$(4yz^3 + 3x^2 z^2 - xye^{xyz}) \frac{\partial z}{\partial x} = yze^{xyz} - 2xz^3.$$

So

$$\frac{\partial z}{\partial x} = \frac{yze^{xyz} - 2xz^3}{4yz^3 + 3x^2 z^2 - xye^{xyz}}.$$

A similar gruesome calculation gives

$$\frac{\partial z}{\partial y} = \frac{xze^{xyz} - z^4}{4yz^3 + 3x^2 z^2 - xye^{xyz}}$$

**Problem 44.** (a) When the direction vector  $\mathbf{u}$  points in the same direction as the gradient  $\nabla f$ . (b) When the direction vector  $\mathbf{u}$  points in the opposite direction to the gradient. (c) When  $\mathbf{u}$  is perpendicular to the gradient. (d) When  $\cos(\theta) = \frac{1}{2}$ , ie when the angle between  $\mathbf{u}$  and  $\nabla f$  is  $\frac{\pi}{3}$ .

**Problem 46.**  $\nabla f = \langle 2xy + \sqrt{1+z}, x^2, -\frac{x}{2\sqrt{1+z}} \rangle = \langle 6, 1, \frac{1}{4} \rangle$  at  $(1, 2, 3)$ . The unit vector in the given direction is  $\frac{1}{3} \langle 2, 1, -2 \rangle$ . Taking the dot product gives  $D_{\mathbf{u}}f(1, 2, 3) = \frac{25}{6}$ .

## 2 Solutions to additional problems

**Problem A.** Write  $\alpha(t) = (x(t), y(t))$ , and suppose we are at the point  $(x_0, y_0) = \alpha(t_0)$  on the graph. Note that we have the equation  $F(x(t), y(t)) = 0$ . Differentiating with respect to  $t$ , and applying the chain rule gives  $F_x(x_0, y_0)x'(t_0) + F_y(x_0, y_0)y'(t_0) = 0$ . In other words, the vectors  $\langle F_x(x_0, y_0), F_y(x_0, y_0) \rangle$  and  $\langle x'(t_0), y'(t_0) \rangle$  are orthogonal, so slopes of lines that these vectors span are inverse reciprocals. (I.e one is  $-1/\text{slope of the other}$ .) Now the slope of the tangent line which is parallel to  $\alpha'(t_0)$ , is given by  $y'(t_0)/x'(t_0)$ , and the slope of the line parallel to

$\langle \nabla F(x_0, y_0), \nabla F(x_0, y_0) \rangle$  is  $F_y(x_0, y_0)/F_x(x_0, y_0)$ . So  $y'(t_0)/x'(t_0) = -F_x(x_0, y_0)/F_y(x_0, y_0)$ .

**Problem B.**  $f(0, 0) = a_0$ ,  $f_x(0, 0) = a$ , and  $f_y(0, 0) = b$ . The linearization is therefore given by  $a_0 + ax + by$ . My guess is that the linearization at  $(0, 0)$  of a polynomial of high degree is just the constant and the linear terms.

**Problem C.** Why not just let  $g(x) = F(x, 2) = 8 \sin(2x) + e^{2x}$ ?

**Problem D.** As suggested in the hint, there's no  $\theta$  in this equation. That means that the surface looks the same in every direction (as measured by  $\theta$ ) or in other words that it has circular symmetry around the  $z$ -axis.

**Problem E.** There is no such function. For the directional derivative is given by  $\nabla f(x_0, y_0) \cdot \mathbf{u}$ , and if this is negative for one value of  $\mathbf{u}$ , it will be positive for the opposite direction given by  $-\mathbf{u}$ .

**Problem F.** We can discuss this at the review.

**Problem G.** There are many possible solutions to this problem; here's one. Start with  $f(x, y) = \sqrt{x^2 + y^2}$ , whose contour line at height  $k$  is the circle of radius  $k$ . We want the same contours for all whole numbers  $k$ , but different contours for other  $k$ . For instance  $g(x, y) = \cos(2\pi\sqrt{x^2 + y^2})\sqrt{x^2 + y^2}$  will work, since  $\cos(2\pi\sqrt{x^2 + y^2})$  is 1 whenever  $\sqrt{x^2 + y^2}$  is a whole number, and less than 1 otherwise.

**Problem H.** The main observation is that any function written as a sum of other functions, say  $f(x, y) = g(x, y) + h(x, y)$ , all of the derivatives of  $f$  are sums of the corresponding derivatives of  $g$  and  $h$ . In particular,  $f_{xy} = g_{xy} + h_{xy}$  and  $f_{yx} = g_{yx} + h_{yx}$ . So if Clairaut's theorem holds for  $g$  and  $h$ , it must hold for  $f$ . Similarly, if Clairaut's theorem holds for a function  $f$ , it holds for  $cf$  where  $c$  is any constant. Since any polynomial is a sum of terms of the form  $cx^m y^n$ , Clairaut's theorem for polynomials will follow once we've proven it for terms of the form  $x^m y^n$ . We do that by direct calculation:  $\frac{\partial^2}{\partial x \partial y}(x^m y^n) = mn x^{m-1} y^{n-1} = \frac{\partial^2}{\partial y \partial x}(x^m y^n)$ .